

Linear Regression Part 2

DATA 606 - Statistics & Probability for Data Analytics

Jason Bryer, Ph.D. and Angela Lui, Ph.D.

April 19, 2023

Announcements

Project Presentations

Please take a look at the sign-up sheet even if you already signed up. We had to move the two Thursday evening time slots to Monday evening.

You can sign up for a presentation slot on this Google Sheet:

<https://docs.google.com/spreadsheets/d/1xcWRZuM4OO0ePR7rqO43tuXVGLQnc0asK1tODdEqfo/edit?usp=sharing>

Example project slides located here:

<https://github.com/jbryer/DATA606Spring2023/tree/main/Project>



One Minute Paper Results

What was the most important thing you learned during this class?

regression
linear
two
can
equation based
understanding squared
difference using
actual line
best relationship
squares
residual idea predicted
error representation
covariance model
correlation

What important question remains unanswered for you?

multivariate
distributed residuals
true can effect
sat get example
linear model regression
class squared normally



NYS Report Card

NYS publishes data for each school in the state. We will look at the grade 8 math scores for 2012 and 2013. 2013 was the first year the tests were aligned with the Common Core Standards. There was a lot of press about how the passing rates for most schools dropped. Two questions we wish to answer:

1. Did the passing rates drop in a predictable manner?
2. Were the drops different for charter and public schools?

```
load('../course_data/NYSReportCard-Grade7Math.Rda')
names(reportCard)
```

```
## [1] "BEDSCODE"      "School"        "NumTested2012"  "Mean2012"
## [5] "Pass2012"       "Charter"        "GradeSubject"   "County"
## [9] "BOCES"          "NumTested2013"  "Mean2013"       "Pass2013"
```



reportCard Data Frame

Show 3 entries

Search:

BEDSCODE	School	NumTested2012	Mean2012	Pass2012	Charter	GradeSubject	County	BOCES	NumTested2013	Mean2013	Pa
010100010020	NORTH ALBANY ACADEMY	47	649	13	false	Grade 7 Math	Albany	BOCES ALBANY- SCHOH- SCHEECTADY- SARAT	45	268	
010100010030	WILLIAM S HACKETT MIDDLE SCHOOL	212	652	30	false	Grade 7 Math	Albany	BOCES ALBANY- SCHOH- SCHEECTADY- SARAT	250	279	
010100010045	STEPHEN AND HARRIET MYERS MIDDLE SCHOOL	262	670	50	false	Grade 7 Math	Albany	BOCES ALBANY- SCHOH- SCHEECTADY- SARAT	256	284	

Showing 1 to 3 of 1,362 entries

Previous

1 2 3 4 5 ... 454

Next



Descriptive Statistics

```
summary(reportCard$Pass2012)
```

```
##    Min. 1st Qu. Median   Mean 3rd Qu.   Max.  
##    0.00    46.00   65.00   61.73   80.00 100.00
```


```
summary(reportCard$Pass2013)
```

```
##    Min. 1st Qu. Median   Mean 3rd Qu.   Max.  
##    0.00     7.00   20.00   22.83   33.00  99.00
```



Histograms


```
melted <- melt(reportCard[,c('Pass2012', 'Pass2013')])  
ggplot(melted, aes(x=value)) + geom_histogram() + facet_wrap(~ variable, ncol=1)
```



Log Transformation


Since the distribution of the 2013 passing rates is skewed, we can log transform that variable to get a more reasonably normal distribution.

```
reportCard$LogPass2013 <- log(reportCard$Pass2013 + 1)  
ggplot(reportCard, aes(x=LogPass2013)) + geom_histogram(binwidth=0.5)
```



Scatter Plot

```
ggplot(reportCard, aes(x=Pass2012, y=Pass2013, color=Charter)) +  
  geom_point(alpha=0.5) + coord_equal() + ylim(c(0,100)) + xlim(c(0,100))
```



Scatter Plot (log transform)

```
ggplot(reportCard, aes(x=Pass2012, y=LogPass2013, color=Charter)) +  
  geom_point(alpha=0.5) + xlim(c(0,100)) + ylim(c(0, log(101)))
```



Correlation

```
cor.test(reportCard$Pass2012, reportCard$Pass2013)
```

```
##  
##      Pearson's product-moment correlation  
##  
## data: reportCard$Pass2012 and reportCard$Pass2013  
## t = 47.166, df = 1360, p-value < 2.2e-16  
## alternative hypothesis: true correlation is not equal to 0  
## 95 percent confidence interval:  
##  0.7667526 0.8071276  
## sample estimates:  
##      cor  
## 0.7877848
```



Correlation (log transform)

```
cor.test(reportCard$Pass2012, reportCard$LogPass2013)
```

```
##  
##      Pearson's product-moment correlation  
##  
## data: reportCard$Pass2012 and reportCard$LogPass2013  
## t = 56.499, df = 1360, p-value < 2.2e-16  
## alternative hypothesis: true correlation is not equal to 0  
## 95 percent confidence interval:  
##  0.8207912 0.8525925  
## sample estimates:  
##       cor  
## 0.8373991
```



Linear Regression

```
lm.out <- lm(Pass2013 ~ Pass2012, data=reportCard)
summary(lm.out)
```

```
## 
## Call:
## lm(formula = Pass2013 ~ Pass2012, data = reportCard)
## 
## Residuals:
##     Min      1Q  Median      3Q     Max 
## -35.484  -6.878  -0.478   5.965  51.675 
## 
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)    
## (Intercept) -16.68965    0.89378  -18.67   <2e-16 ***
## Pass2012     0.64014    0.01357   47.17   <2e-16 ***
## ---      
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 
## Residual standard error: 11.49 on 1360 degrees of freedom
## Multiple R-squared:  0.6206,    Adjusted R-squared:  0.6203 
## F-statistic: 2225 on 1 and 1360 DF,  p-value: < 2.2e-16
```



Linear Regression (log transform)

```
lm.log.out <- lm(LogPass2013 ~ Pass2012, data=reportCard)
summary(lm.log.out)
```

```
## 
## Call:
## lm(formula = LogPass2013 ~ Pass2012, data = reportCard)
## 
## Residuals:
##     Min      1Q  Median      3Q     Max 
## -3.3880 -0.2531  0.0776  0.3461  2.7368 
## 
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)    
## (Intercept) 0.307692   0.046030   6.685 3.37e-11 ***
## Pass2012    0.039491   0.000699  56.499 < 2e-16 ***
## ---      
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
## 
## Residual standard error: 0.5915 on 1360 degrees of freedom
## Multiple R-squared:  0.7012,    Adjusted R-squared:  0.701 
## F-statistic: 3192 on 1 and 1360 DF,  p-value: < 2.2e-16
```



Did the passing rates drop in a predictable manner?

Yes! Whether we log transform the data or not, the correlations are statistically significant with regression models with R^2 greater than 62%.


To answer the second question, whether the drops were different for public and charter schools, we'll look at the residuals.

```
reportCard$residuals <- resid(lm.out)
reportCard$residualsLog <- resid(lm.log.out)
```




Distribution of Residuals

```
ggplot(reportCard, aes(x=residuals, color=Charter)) + geom_density()
```



Distribution of Residuals

```
ggplot(reportCard, aes(x=residualsLog, color=Charter)) + geom_density()
```



Null Hypothesis Testing

H_0 : There is no difference in the residuals between charter and public schools.

H_A : There is a difference in the residuals between charter and public schools.

```
t.test(residuals ~ Charter, data=reportCard)
```

```
##  
##      Welch Two Sample t-test  
##  
## data: residuals by Charter  
## t = 6.5751, df = 77.633, p-value = 5.091e-09  
## alternative hypothesis: true difference in means between group FALSE and group TRUE is not equal to 0  
## 95 percent confidence interval:  
##   6.411064 11.980002  
## sample estimates:  
## mean in group FALSE mean in group TRUE  
##             0.479356          -8.716177
```



Null Hypothesis Testing (log transform)

```
t.test(residualsLog ~ Charter, data=reportCard)
```

```
##  
##      Welch Two Sample t-test  
##  
## data: residualsLog by Charter  
## t = 4.7957, df = 74.136, p-value = 8.161e-06  
## alternative hypothesis: true difference in means between group FALSE and group TRUE is not equal to 0  
## 95 percent confidence interval:  
##  0.2642811 0.6399761  
## sample estimates:  
## mean in group FALSE  mean in group TRUE  
##                 0.02356911        -0.42855946
```



Polynomial Models (e.g. Quadratic)

It is possible to fit quadratic models fairly easily in R, say of the following form:

$$y = b_1x^2 + b_2x + b_0$$

```
quad.out <- lm(Pass2013 ~ I(Pass2012^2) + Pass2012, data=reportCard)
summary(quad.out)$r.squared
```

```
## [1] 0.7065206
```

```
summary(lm.out)$r.squared
```

```
## [1] 0.6206049
```



Quadratic Model


```
summary(quad.out)
```

```
##  
## Call:  
## lm(formula = Pass2013 ~ I(Pass2012^2) + Pass2012, data = reportCard)  
##  
## Residuals:  
##      Min       1Q   Median       3Q      Max  
## -46.258  -4.906  -0.507   5.430  43.509  
##  
## Coefficients:  
##                 Estimate Std. Error t value Pr(>|t|)  
## (Intercept) 7.0466153 1.4263773  4.940 8.77e-07 ***  
## I(Pass2012^2) 0.0092937 0.0004659 19.946 < 2e-16 ***  
## Pass2012     -0.3972481 0.0533631 -7.444 1.72e-13 ***  
## ---  
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## Residual standard error: 10.11 on 1359 degrees of freedom  
## Multiple R-squared: 0.7065, Adjusted R-squared: 0.7061  
## F-statistic: 1636 on 2 and 1359 DF, p-value: < 2.2e-16
```



Scatter Plot


```
ggplot(reportCard, aes(x=Pass2012, y=Pass2013)) + geom_point(alpha=0.2) +  
  geom_smooth(method='lm', formula=y ~ x, size=2, se=FALSE) +  
  geom_smooth(method='lm', formula=y ~ I(x^2) + x, size=2, se=FALSE) +  
  coord_equal() + ylim(c(0,100)) + xlim(c(0,100))
```



Let's go crazy, cubic!

```
cube.out <- lm(Pass2013 ~ I(Pass2012^3) + I(Pass2012^2) + Pass2012, data=reportCard)  
summary(cube.out)$r.squared
```


```
## [1] 0.7168206
```



Be careful of overfitting...

OVERFITTING


Overfitting occurs when a model starts to memorize the aspects of the training set and in turn loses the ability to generalize




Chris Albon

Source: Chris Albon [@chrisalbon](https://MachineLearningFlashCards.com)


OVERFIT VS UNDERFIT



OVERFIT
"HIGH VARIANCE"



IDEAL




UNDERFIT
"HIGH BIAS"



Loess Regression

```
ggplot(reportCard, aes(x=Pass2012, y=Pass2013)) + geom_point(alpha=0.2) +  
  geom_smooth(method='lm', formula=y~poly(x,2,raw=TRUE), size=2, se=FALSE) +  
  geom_smooth(method='loess', formula = y ~ x, size=2, se=FALSE, color = 'purple') +  
  coord_equal() + ylim(c(0,100)) + xlim(c(0,100))
```



```
library('VisualStats')  
library('ShinyDemo')  
shiny_demo('loess', package = 'VisualStats')
```

See this site for more info:

<https://jbryer.github.io/VisualStats/articles/loess>



Shiny App

```
shiny::runGitHub('NYSchools','jbryer',subdir='NYSReportCard')
```

See also the Github repository for more information: <https://github.com/jbryer/NYSchools>



One Minute Paper

Complete the one minute paper:

<https://forms.gle/p9xcKcTbGiyYSz368>

1. What was the most important thing you learned during this class?
2. What important question remains unanswered for you?

